

Epidemiology of Acute Myocarditis/Pericarditis in Hong Kong Adolescents Following Comirnaty Vaccination.

Gilbert T Chua^{*1,2}, FHKCPaed, FHKAM(Paed), Mike Yat Wah Kwan^{*3}, MSc(Applied Epidemiology) CUHK, Celine SL Chui^{4,5,6}, PhD, Robert David Smith⁴, PhD, Edmund Chi-Lok Cheung⁷, BSc, Tian Tian Ma⁶, PhD, Miriam TY Leung^{6,7}, BPharm, Sabrina Siu Ling Tsao^{1,2}, MBBS(UK), FACC(USA), Elaine Kan⁸, FRCR, FHKCR, Wing Kei Carol Ng⁸, FRCR, FHKAM, Victor Chi Man Chan⁹, MRCPCH, Shuk Mui Tai⁹, MBChB, Tak Ching Yu⁹, MRCPCH, Kwok Piu Lee⁹, FHKAM, FHKCPaed, Joshua Sung Chih Wong³, FHKCPaed, FHKAM(Paed), Ying Kit Lin³, MRCPCH, Chi Chiu Shek³, FHKCPaed, FRCPCH, Agnes Sze Yin Leung¹⁰, FHKCPaed, FHKAM(Paed), Chit Kwong Chow¹¹, FHKCPaed, FHKAM(Paed), Ka Wah Li¹², FHKCPaed, FHKAM(Paed), Johnny Ma^{13,14,15,16}, MMed, Wai Yuk Fung^{13,14,15,16}, MBChB, FRCR, Daniel Lee¹⁷, FRCR, Ming Yen Ng^{18,19}, FSCMR, FACC, Wilfred Hing Sang Wong¹, PhD, Hing Wai Tsang¹, PhD, Janette Kwok²⁰, PhD, FRCPA, Daniel Leung¹, PhD (candidate), Kin Lai Chung²¹, MBBS, FRCS, Chun Bong Chow¹, MD, FRCPCH, Godfrey Chi Fung Chan^{1,2}, MD, FRCPCH(UK), Wing Hang Leung^{1,2}, MBBS(HK), PhD(Johns Hopkins), Kelvin Kai Wang To²², MD, FRCPath, Kwok Yung Yuen²², MD, FRCPath, Yu Lung Lau^{1,2}, MD(Hon), FRCPCH, Ian Chi Kei Wong^{6,7,23}, PhD, FRCPCH(Hon), Patrick Ip¹, MPH, FRCPCH(UK)

*Equal Contributions

1. Department of Paediatrics and Adolescent Medicine, Li Ka Shing Faculty of Medicine,
The University of Hong Kong, Hong Kong SAR, China
2. Department of Paediatrics, Hong Kong Children's Hospital
3. Department of Paediatrics and Adolescent Medicine, Princess Margaret Hospital,
Hong Kong SAR, China
4. School of Nursing, The University of Hong Kong, Hong Kong SAR, China
5. School of Public Health, The University of Hong Kong, Hong Kong SAR, China
6. Laboratory of Data Discovery for Health (D24H), Hong Kong Science and Technology
Park, Hong Kong SAR, China
7. Centre for Safe Medication Practice and Research, Department of Pharmacology and
Pharmacy, The University of Hong Kong, Hong Kong SAR, China
8. Department of Radiology, Hong Kong Children's Hospital
9. Department of Paediatrics and Adolescent Medicine, Pamela Youde Nethersole
Eastern Hospital, Hong Kong SAR, China
10. Department of Pediatrics, Prince of Wales Hospital, The Chinese University of Hong
Kong, Hong Kong SAR, China

- 11.** Department of Paediatrics and Adolescent Medicine, United Christian Hospital, Hong Kong SAR, China
- 12.** Department of Paediatrics and Adolescent Medicine, Tuen Mun Hospital, Hong Kong SAR, China
- 13.** Department of Radiology, Caritas Medical Centre, Hong Kong SAR, China
- 14.** Department of Radiology, North Landau Hospital, Hong Kong SAR, China
- 15.** Department of Radiology, Princess Margaret Hospital, Hong Kong SAR, China
- 16.** Department of Radiology, Yan Chai Hospital, Hong Kong SAR, China
- 17.** Department of Radiology, Pamela Youde Nethersole Eastern Hospital, Hong Kong SAR, China
- 18.** Department of Diagnostic Radiology, The University of Hong Kong
- 19.** Department of Medical Imaging, The University of Hong Kong-Shenzhen Hospital, China
- 20.** Division of Transplantation and Immunogenetics, Department of Pathology, Queen Mary Hospital, Hong Kong SAR, China
- 21.** Quality & Safety Division, Hospital Authority Head office, Hong Kong SAR, China
- 22.** Department of Microbiology, Carol Yu Centre for Infection, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong SAR, China
- 23.** Research Department of Practice and Policy, UCL School of Pharmacy, University College, London, United Kingdom

Corresponding Author

Dr Patrick Ip, Clinical Associate Professor

Department of Paediatrics and Adolescent Medicine

Li Ka Shing Faculty of Medicine, The University of Hong Kong

Hong Kong SAR, China

Email: patricip@hku.hk

Article Summary:

There is an increased risk of acute myocarditis/pericarditis in adolescent males following Comirnaty vaccination, especially following the second dose, providing the direct evidence to support the change in recommending Hong Kong's adolescents to receive one dose of the Comirnaty vaccine.

Abstract

Background

Age-specific incidence of acute myocarditis/pericarditis in adolescents following Comirnaty vaccination in Asia is lacking. This study aimed to study the clinical characteristics and incidence of acute myocarditis/pericarditis among Hong Kong adolescents following Comirnaty vaccination.

Methods:

This is a population cohort study in Hong Kong that monitored adverse events following immunization through a pharmacovigilance system for COVID-19 vaccines. All adolescents aged between 12 and 17 years following Comirnaty vaccination were monitored under the COVID-19 vaccine Adverse Event Response and Evaluation Programme. The clinical characteristics and overall incidence of acute myocarditis/pericarditis in adolescents following Comirnaty vaccination were analysed.

Results:

Between 14 June 2021 and 4 September 2021, 33 Chinese adolescents who developed acute myocarditis/pericarditis following Comirnaty vaccination were identified. 29 (87.88%) were males and 4 (12.12%) were females, with a median age of 15.25 years. 27 (81.82%) and 6 (18.18%) cases developed acute myocarditis/pericarditis after receiving the second and first dose, respectively. All cases are mild and required only conservative management.

The overall incidence of acute myocarditis/pericarditis was 18.52 (95% Confidence Interval [CI], 11.67-29.01) per 100,000 persons vaccinated. The incidence after the first and second doses were 3.37 (95%CI 1.12-9.51) and 21.22 (95%CI 13.78-32.28 per 100,000 persons vaccinated, respectively. Among male adolescents, the incidence after the first and second doses were 5.57 (95% CI 2.38-12.53) and 37.32 (95% CI 26.98-51.25) per 100,000 persons vaccinated.

Conclusions:

There is a significant increase in the risk of acute myocarditis/pericarditis following Comirnaty vaccination among Chinese male adolescents, especially after the second dose.

Keywords: Myocarditis; Pericarditis; adolescents; Comirnaty; Hong Kong

Accepted Manuscript

Introduction

The COVID-19 infection in children is generally mild, but serious complications, such as Paediatric Multisystem Inflammatory Syndrome – Temporally Associated with SARS-CoV-2 (PIMS-TS), can occur.¹ Prolonged social distancing policies have also led to significant psychosocial impacts on children and their families in the community.² Enormous efforts have been made to control the spread of the virus through universal vaccination to achieve herd immunity to return us to a semblance of normality.

Currently, the Vaccination Programme of the Hong Kong Government has authorized two COVID-19 vaccines: the CoronaVac from Sinovac Biotech (Hong Kong) Limited and Comirnaty vaccine (BNT162b2) from Fosun-BioNTech. On 14 June 2021, the Government of the Hong Kong Special Administrative Region (HKSAR) commenced vaccination of the Comirnaty vaccine (BNT162b2) from Fosun-BioNTech to adolescents by lowering of the age limit from 16 to 12 years after reviewing the available evidence by the Advisory Panel on COVID-19 Vaccines of the Food and Health Bureau, HKSAR Government.³ The Drug Office of the Department of Health (DH), the drug regulatory authority in Hong Kong, has implemented a pharmacovigilance system for COVID-19 vaccines that monitors reports of adverse events following immunization (AEFI). The COVID-19 vaccine Adverse event Response and Evaluation (CARE) Programme was set up, an active surveillance system, to evaluate AEFI data from the general population using electronic medical records from Hospital Authority and vaccination records from the DH. The CARE programme actively identifies AEFI and conduct epidemiological study to evaluate the association between vaccinations and subsequent adverse event.^{4, 5}

The Comirnaty is a messenger RNA (mRNA) vaccine that is highly effective in preventing hospitalizations and deaths due to COVID-19.⁶ Although Comirnaty has a favourable safety profile, various regulatory agencies have advocated continuous monitoring of its safety, as rare and long-term adverse reactions might not have been detected in the clinical trials and early post-marketing reports.⁷ Recently, there have been emerging case reports of acute myocarditis following mRNA COVID-19 vaccination in healthy young adolescent and adult males.⁸⁻¹⁰ The United Kingdom has only approved offering one dose of the Pfizer-BioNTech vaccine to healthy adolescents aged 12 to 15 years old so far, instead of giving the recommended two doses.¹¹ Yet an in-depth population-based investigation of the age-specific incidence of acute myocarditis/pericarditis following mRNA COVID-19 vaccination in Asian adolescents is lacking. This study aims to report the clinical characteristics and estimate the incidence of acute myocarditis following vaccination with Comirnaty in adolescents in Hong Kong.

Methods

This was a population cohort study aimed at identifying all suspected cases of acute myocarditis in adolescents aged between 12 and 17 years who received the Comirnaty vaccine between 14 June 2021 and 4 September 2021. All individuals receiving the Comirnaty vaccine have also consented to their vaccination records being linked to their corresponding comprehensive electronic health records held by the Hospital Authority (HA), the major publicly funded healthcare provider, through the CARE Programme.⁴ All suspected cases of acute myocarditis/pericarditis that occur within 14 days after receiving either the first or the second dose of the Comirnaty vaccine and admitted to one of the HA hospitals are reported to the Advanced Incident Reporting System (AIRS) on admission, a system for HA to report adverse drug events and AEFI to DH.

Suspected cases of acute myocarditis/pericarditis who received Comirnaty vaccines during the study period were investigated according to the Hong Kong Paediatric Investigation Protocol for Comirnaty-related Myocarditis/Pericarditis (Supplementary file 1), which was implemented in all HA hospitals. Demographics including date of birth, sex, ethnicity, date of receiving the first and the second dose of COVID-19 vaccines, symptoms, date of onset, and past medical histories were reviewed. Microbiological investigations including nasopharyngeal swab (NPS) for SARS-CoV-2 and common respiratory viruses including influenza A/B/C, parainfluenza virus 1/2/3/4, adenovirus, human metapneumovirus, and respiratory syncytial virus, and throat and rectal swabs for enteroviruses were tested. SARS-CoV-2 anti-receptor binding domain (RBD) and anti-nucleocapsid protein (NP) antibodies were tested to differentiate whether the patients had a history of COVID-19 infection. Cardiac enzymes, including high-sensitivity troponin I (hsTnI), high-sensitivity troponin T (hsTnT or TnT), electrocardiogram (ECG), and echocardiogram were serially monitored. ECGs were interpreted by one single investigator (SST). Echocardiograms were performed and interpreted by the cardiologists of each admitting hospital. Cardiac magnetic resonance imaging (cMRI) was performed within two weeks of symptoms onset either at the admission hospital, or referred to the Hong Kong Children's Hospital if no slots were immediately available. cMRI images were interpreted by the radiologists of each MRI unit. The study team followed the myocarditis and pericarditis case definitions created by the Cardiovascular Injury-Coalition for Epidemic Preparedness Innovations (CEPI) and the Brighton Working Group.¹²

Estimation of Incidence and Statistical Analysis

Vaccination records within the study period were extracted from the DH in Hong Kong since the commencement of mass COVID-19 vaccinations in adolescents aged 12 to 17 years on 14 June 2021 to 4 September 2021. The cut-off date for follow-up time was 18 September 2021, allowing for all participants to have a 14-day follow-up period. De-identified electronic health records were retrieved from the HA Clinical Data Analysis and Reporting System (CDARS), which has been successfully used in a previous COVID-19 vaccine-related pharmacovigilance study.⁴ Subjects with a history of primary myocarditis/pericarditis prior to the study period were excluded. Cases of acute myocarditis/pericarditis following Comirnaty vaccination were identified if they occurred within 14 days of either the first or the second vaccine dose. We estimated the background rate of acute myocarditis/pericarditis, cases of the first primary diagnosis were extracted from CDARS from 2011 to 2020 using data available from 14 June to the 4 September in each year. For each year, those with a history of acute myocarditis/pericarditis in the prior year to the study period were censored.

Separated cases related to the first dose or to the second dose were also calculated. Acute myocarditis/pericarditis related to the first dose was defined as the first cases within 14 days of the first dose. Acute myocarditis/pericarditis related to the second dose was defined as the first cases within 14 days of the second dose. The 14 days was the upper end of the reporting of myocarditis/pericarditis cases following vaccination according to the DH and HA reporting policies. The incidence of clinically confirmed myocarditis/pericarditis per 100,000 doses administered as well as number of cases per 100,000 doses for first dose and second dose were estimated. We calculated 95% confidence intervals (95% CI) for all incidences calculated using Poisson distribution. The incidence rate of acute myocarditis/pericarditis associated with the Comirnaty vaccine was

compared with the background incidence rate of acute myocarditis/pericarditis in 2020 using per 100,000 per 14-days. Sensitivity analyses were conducted using (1) the background incidence rate in 2018 and 2019 and the average background incidence rate from 2011 to 2020 using per 100,000 per 14-days and (2) changed the incidence using per 28-days. Subgroup analysis was conducted by sex. Some comparisons to background years were not possible as there were zero cases of myocarditis/pericarditis recorded in background years. Median and interquartile ranges (IQR) were used to describe skewed data. All statistical tests were two-sided and p-values at a level of 5% were considered statistically significant. Statistical analyses were conducted using R version 4.0.3 (www.R-project.org). For quality assurance, two investigators (ECC and RDS) independently conducted the statistical analyses.

Ethical approval

This study was approved by the Institutional Review Board of the University of Hong Kong/Hospital Authority Hong Kong West Cluster (UW21-149 and UW21-138) and the Department of Health Ethics Committee (LM21/2021).

Results

Between 14 June and 4 September 2021, a total of 33 cases of myocarditis/pericarditis within 14 days following vaccination with Comirnaty were identified. Twenty-five (75.76%) were definite, 7 (21.21%) were probable and 1 (3.03%) were possible cases (Table 1). The patients were all Chinese adolescents with no history of cardiac diseases. 29 (87.88%) were males and 4 (12.12%) were females, with a median age of 15.25 years. 27 (81.82%) and 6 (18.18%) cases developed acute myocarditis/pericarditis after receiving the second and first dose, respectively. These patients developed myocarditis/pericarditis at a median of 2 days after receiving the last dose of the vaccine.

All of them presented with chest pain. Three cases (9.09%) had normal troponin levels, two of them were cases of definite pericarditis and one possible myocarditis. 6 (18.18%) had normal ECGs, 25 (75.76%) had normal echocardiograms, and 7 (21.88%) had normal cMRI. None had significant arrhythmias. All patients had no identifiable infections. They also had no current and past history of COVID-19 infection, as evidenced by a negative SARS-CoV-2 PCR on admission and the absent of anti-SARS-CoV-2 NP antibodies in their serum. All patients had mild diseases requiring no treatment or symptomatic relief by non-steroidal anti-inflammatory drugs (NSAIDs). They spontaneously recovered without the need of systemic steroids, intravenous immunoglobulins, inotropic or circulatory support.

There have been 305,406 doses of Comirnaty vaccine administered to 178,163 individuals aged 12-17 years (88,357 [49.59%] are female) since the commencement of the vaccination program on 14 June 2021 until 4 September 2021. This represented 51.84% of the population between 12 and 17 years (178,163/343,700) in Hong Kong in mid-2021.¹³ The overall incidence for acute myocarditis/pericarditis was 18.52 (95% CI 11.67-29.09) per 100,000 persons vaccinated. The incidence after the first and second doses were 3.37 (95%CI 1.12-9.51) and 21.22 (95%CI 13.78-32.28) per 100,000 persons vaccinated, respectively (Table 2). Incidence was higher among male adolescents compared to females (Table 2). Incidence rates compared with previous years' background rates are shown in Table 2 and Supplementary Tables 1 – 3. Compared to the background incidence rate of acute myocarditis/pericarditis in 2020 there were significantly higher incidence rate differences in those vaccinated (Table 2). Sensitivity analyses using the background incidence rate in 2018, 2019 and 2020 and the average background incidence rate from 2011 to 2020 using per 100,000 per 28-days also demonstrated significantly higher incidence rate differences in those vaccinated which was consistent with the main results (Supplementary tables 4 – 8).

Among males after their first dose, there was a significantly higher incidence rate difference compared the background rate in 2020. After the second dose there was significantly higher incidence rate difference between the background rate in 2020 and all participant and males (Tables 3).

Discussion

To our best knowledge, this is the first study in adolescents using data from the territory-wide post-COVID-19 vaccination monitoring system to analyse the incidence of acute myocarditis/pericarditis associated with the Comirnaty vaccine for adolescents in Asia.

Our analysis revealed that the overall incidence of acute myocarditis/pericarditis in adolescent following the Comirnaty vaccination was 18.52 per 100,000 persons vaccinated. Majority cases involved healthy adolescent males after receiving the second dose. No other infective causes including SARS-CoV-2 infection were identified. Conservative management with NSAIDs was sufficient. This higher incidence of myocarditis/pericarditis following Comirnaty vaccination than other jurisdictions is likely related to the heightened vigilance of healthcare professionals and the public,¹⁴ as well as the highly efficient CARE Programme for the monitoring and reporting of AEFI across Hong Kong.¹⁰ Our pharmacovigilance system was able to capture mild cases of acute myocarditis/pericarditis and reveal the real-world incidence of acute myocarditis/pericarditis following the Comirnaty vaccination. Since the Pfizer-BioNTech vaccine was approved for large-scale immunization in many countries, there has been a higher observed risk of acute myocarditis/pericarditis among younger males receiving this vaccine.¹⁵ The first reports in Israel were of five young males who developed mild myocarditis following vaccination with the BioNTech mRNA COVID-19 vaccine.¹⁶ Subsequently, 23 US military males reported developing myocarditis

after administering more than 2.8 million doses of either the Moderna or BioNTech mRNA COVID-19 vaccines to military personnel.⁹ In children, so far, only one case series reported myocarditis following vaccinations with mRNA COVID-19 vaccines. These seven cases were males aged 14 to 19 years who presented with transient mild symptoms, elevated troponin, and MRI changes suggestive of acute myocarditis or perimyocarditis. They were treated with NSAIDs, steroids, or intravenous immunoglobulin.⁸ So far, all adults and adolescents with myocarditis/pericarditis following COVID-19 vaccinations, including those reported in the current study, have been mild cases.¹⁷ While the pathophysiology of acute myocarditis/pericarditis following the mRNA COVID-19 vaccine is still unclear and the observation that only mRNA-based COVID-19 vaccines are associated with acute myocarditis remains unexplained. The causal association between mRNA vaccine and myopericarditis has recently been suggested in a mouse model. Higher systemic levels of mRNA lipid nanoparticles due to inadvertent intravenous injection or rapid return from the lymphatic circulation was proposed to increase this risk.¹⁸ Further studies to delineate the pathophysiology of acute myocarditis/pericarditis associated with mRNA-based COVID-19 vaccines is urgently needed.

The US Center for Disease Control and Prevention (CDC) reported that the expected rates of myocarditis/pericarditis following the Comirnaty vaccination would be the highest among males aged between 12 and 29 years old, estimating 40.6 per million second doses administered.¹⁰ The incidence rate of myocarditis/pericarditis following the Comirnaty vaccination in Hong Kong was much higher than those reported from the United States.^{10, 19} However, it is important to note that the risk of myocardial injury in healthy young individuals including athletes following COVID-19 infection is also considerably high,²⁰ ranging from asymptomatic cases with abnormal cMRI only to fulminant myocarditis due to COVID-19.^{21, 22} Preliminary data in Israel demonstrated a 51% effectiveness after receiving one dose Pfizer-BioNTech vaccine among older adults.²³ As there have been essentially no local transmission of SARS-CoV-2 in Hong Kong since May 2021,²⁴ balancing the

risk of acute myocarditis/pericarditis after receiving the second dose and the benefit of vaccination to protect complications related to COVID-19 infection, the Scientific Committee on Vaccine Preventable Diseases and the Scientific Committee on Emerging and Zoonotic Diseases under the Centre for Health Protection of the Department of Health of Hong Kong recommended adolescents between 12 and 17 years to receive one dose of the Comirnaty vaccine, instead of two doses, on 15 September 2021.²⁵ While our study provided the most comprehensive epidemiology of myocarditis/pericarditis following Comirnaty vaccination before the policy change, ongoing observations on the incidence of myocarditis/pericarditis following the Comirnaty vaccination with one-dose Comirnaty vaccination as well as the rate of COVID-19 infections among adolescents in Hong Kong shall be conducted to provide real-world evidence on the risk and benefit of the policy change.

This study has several strengths and limitations. All subjects presented to the accident and emergency department or in the outpatient clinics in the public system received comprehensive reviews and investigations to rule out the possibility of myocarditis/pericarditis because of viral infection, and cMRI to confirm subtle inflammation of the myocardium. However, asymptomatic subjects and subjects with transient and subtle symptoms of acute myocarditis/pericarditis, such as tachycardia and mild chest discomfort, might not seek medical consultation or have sought medical consultation in the private sector which were not reported. Some patients had negative MRI results because not all MRI suites in Hong Kong's public hospitals have the capability for T1 and T2 mapping to calculate the extracellular volume, leading to lower sensitivities and unable to meet the 2018 Lake Louise Criteria for the diagnosis of myocarditis. Furthermore, the incidence of acute myocarditis/pericarditis following the COVID-19 vaccination remained to be high, possibly attributed to increased awareness of possible acute myocarditis/pericarditis following vaccination with COVID-19 vaccines compared with other jurisdictions, as well as to the CARE Programme to capture AEFI. The high incidence of acute myocarditis/pericarditis following Comirnaty vaccination among adolescents presented in this study is representable as the HA receives majority of emergency

admissions in Hong Kong.⁴ Finally, different criteria were likely used by clinicians in generating a diagnostic code among the non-vaccinated individuals for the calculation of the background myocarditis/pericarditis incidence as it was in a non-research setting. Nevertheless, we have included myocarditis and pericarditis of all causes, including idiopathic cases, for the calculation of the background incidence.

Conclusion

Chinese adolescent males have a higher risk of acute myocarditis/pericarditis following vaccination with Comirnaty, especially after the second dose. Medical professionals and recipients of the Comirnaty vaccine should be vigilant regarding the symptoms of acute myocarditis/pericarditis. Observations on the incidence of myocarditis/pericarditis following the Comirnaty vaccination after changing to one-dose vaccination as well as the rate of COVID-19 infections among adolescents shall be conducted.

Acknowledgements:

We thank the Drug Office of the Department of Health and the Hospital Authority for providing vaccination and clinical data. We also thank Dr Bernard Chan, Wan-Mui Chan, Allen Wing-Ho Chu, and Jonathan Daniel Ip for their technical support and administrative assistance.

Author contributions:

Dr Patrick Ip and Prof Ian Wong have assessed and verified the data.

Concept and design: Dr Mike Kwan, Prof YL Lau, Prof Ian Wong, and Dr Patrick Ip

Acquisition of data: Dr Mike Kwan, Dr Joshua Wong, Dr Victor Chan, Dr Shuk Mui Tai, Dr Kwok Piu Lee, Dr Ying Kit Lin, Dr Chi Chiu Shek, Dr Sabrina Tsao, Dr Elaine Kan, Dr Johnny Ma, Dr Daniel Lee, Dr Janette Kwok, Dr Kelvin To, Prof Godfrey Chan, Dr Hing Wai Tsang, Dr Agnes Leung, Dr Ming Yen Ng, Dr Chit Kwong Chow, Dr Ka Wah Li

Statistical analysis: Dr Celine Chui, Dr Wilfred Wong, Mr Edmund Cheung, Dr Tian Tian Ma and Dr Robert Smith

Interpretation of data: All authors.

Literature review: Dr Hing Wai Tsang, Mr Daniel Leung, Ms Miriam Leung, Prof Kwok Yung Yuen, Prof Wing Hang Leung, Dr Kin Lai Chung and Dr Gilbert Chua

Drafting of the manuscript: Dr Gilbert Chua and Dr Mike Kwan

Figures: Dr Robert Smith

Critical revision of the manuscript for important intellectual content: All authors.

Funding/Support:

Research Grant from the Food and Health Bureau, the Government of the Hong Kong Special Administrative Region (Ref. No. COVID19F01). ICKW reports support from Health and Medical Research Fund of Hong Kong Government (Ref: COVID19F01 to The University of Hong Kong).

Role of Funding Source:

This is a regulatory pharmacovigilance study initiated by the Department of Health (DH) and funded by the Food and Health Bureau of the Government of the Hong Kong Special Administrative Region. The corresponding author has full access to all the data in the study and have final responsibility for the decision to submit for publication.

Conflict of interest:

Dr Celine Chui has received grants outside of the submitted work from the Food and Health Bureau of the Hong Kong Government, Hong Kong Research Grants Council, Hong Kong Innovation and Technology Commission, Pfizer, IQVIA, and Amgen; and a personal fee from Primevigilance Ltd.

Dr Agnes Sze Yin Leung received grants outside of the submitted work from the Health and Medical Research Fund, Food and Health Bureau of the Hong Kong Government Special Administrative Region.

Dr Ming Yen Ng have received funding/education grants from the the Food and Health Bureau of the Hong Kong Government, Radiological Society of North America, GE, Lode, Arterys, Bayer, Circle Cardiovascular Imaging and TeraRecon; honoraria for education activities from Boehringer Ingelheim; reports the following leadership roles: Vice Chair of the Education Committee for Society of Cardiovascular Magnetic Resonance and Member of the Corporate Relations Committee for Society of Cardiovascular Computed Tomography.

Prof Godfrey Chi Fung Chan is the CMO of Xellera and advisor of Pangenia.

Prof Yu Lung Lau received Government funding for COVID-19 Vaccinations in Adolescents (COVA), and is the Chairman of the Scientific Committee on Vaccine Preventable Diseases, Centre for Health Protection, HKSAR

Prof Ian Wong has received research funding outside of the submitted work from Amgen, Bristol-Myers Squibb, Pfizer, Janssen, Bayer, GSK, Novartis, Hong Kong Research Grants Council, Hong Kong Health and Medical Research Fund, National Institute for Health Research in England, European Commission, and National Health and Medical Research Council in Australia (Research grants on pharmacoepidemiology to The University of Hong Kong. Outside submitted work); consultancy fee for advising IQVIA on pharmacoepidemiology studies, outside submitted work; payment for expert testimony from Appeal Court in Hong Kong (Expert report on effects of cannabis. Outside submitted work); and speaker fees from Janssen and Medicine in the previous 3 years; reports the following leadership roles: Member of Pharmacy and Poisons Board (This is the regulatory agency in pharmaceutical product licensing), Member of the Expert Committee on Clinical Events Assessment Following COVID-19 Immunization (Advise the Hong Kong Government on safety of COVID-19 vaccines), and Member of the Advisory Panel on COVID-19 Vaccines of the Hong Kong Government (Advise the Hong Kong Government on the emergency use of COVID-19 vaccines). He is also an independent non-executive director of Jacobson Medical in Hong Kong (salaried).

Dr Patrick Ip has received grants outside of the submitted work from the Food and Health Bureau of the Hong Kong Government, Hong Kong Research Grants Council, and Hong Kong Jockey Club Charities Trust.

Miriam Tim Yin Leung reports receiving Honorarium for talk on ADHD.

Wing Kei Carol Ng reports personal honoraria for Guerbet online lecture on paediatric cardiac imaging; holds 100 shares in Moderna stock, 50 shares in Biontech stock since April, owned 100 shares in Pfizer stock from July 2020 to January 2021.

Accepted Manuscript

References

1. Chua GT, Wong JSC, Lam I, et al. Clinical Characteristics and Transmission of COVID-19 in Children and Youths During 3 Waves of Outbreaks in Hong Kong. *JAMA Network Open*. 2021;4(5):e218824-e218824. doi:10.1001/jamanetworkopen.2021.8824
2. Tso WWY, Wong RS, Tung KTS, et al. Vulnerability and resilience in children during the COVID-19 pandemic. *Eur Child Adolesc Psychiatry*. Nov 17 2020:1-16. doi:10.1007/s00787-020-01680-8
3. Persons aged 12 to 15 can make reservations to receive BioNTech vaccine from tomorrow. The Government of the Hong Kong Special Administrative Region. Accessed Jul 22, 2021. <https://www.info.gov.hk/gia/general/202106/10/P2021061000556.htm>
4. Wan EYF, Chui CSL, Lai FTT, et al. Bell's palsy following vaccination with mRNA (BNT162b2) and inactivated (CoronaVac) SARS-CoV-2 vaccines: a case series and nested case-control study. *The Lancet Infectious Diseases*. doi:10.1016/S1473-3099(21)00451-5
5. Li X, Tong X, Yeung WWY, et al. Two-dose COVID-19 vaccination and possible arthritis flare among patients with rheumatoid arthritis in Hong Kong. *Ann Rheum Dis*. Oct 22 2021;doi:10.1136/annrheumdis-2021-221571
6. Polack FP, Thomas SJ, Kitchin N, et al. Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N Engl J Med*. Dec 31 2020;383(27):2603-2615. doi:10.1056/NEJMoa2034577
7. Gee J, Marquez P, Su J, et al. First Month of COVID-19 Vaccine Safety Monitoring - United States, December 14, 2020-January 13, 2021. *MMWR Morb Mortal Wkly Rep*. Feb 26 2021;70(8):283-288. doi:10.15585/mmwr.mm7008e3
8. Marshall M, Ferguson ID, Lewis P, et al. Symptomatic Acute Myocarditis in Seven Adolescents Following Pfizer-BioNTech COVID-19 Vaccination. *Pediatrics*. Jun 4 2021;doi:10.1542/peds.2021-052478
9. Montgomery J, Ryan M, Engler R, et al. Myocarditis Following Immunization With mRNA COVID-19 Vaccines in Members of the US Military. *JAMA Cardiology*. 2021;doi:10.1001/jamacardio.2021.2833
10. Gargano JW, Wallace M, Hadler SC, et al. Use of mRNA COVID-19 Vaccine After Reports of Myocarditis Among Vaccine Recipients: Update from the Advisory Committee on Immunization Practices - United States, June 2021. *MMWR Morb Mortal Wkly Rep*. Jul 9 2021;70(27):977-982. doi:10.15585/mmwr.mm7027e2
11. Young people aged 12 to 15 to be offered a COVID-19 vaccine. GOV.UK. Accessed Sep 19, 2021. <https://www.gov.uk/government/news/young-people-aged-12-to-15-to-be-offered-a-covid-19-vaccine>
12. Myocarditis/Pericarditis Case Definition. Brighton Collaboration. Accessed Jul 22, 2021. <https://brightoncollaboration.us/myocarditis-case-definition-update/>
13. Table 1B : Population by Sex and Age. Census and Statistics Department, The Government of the HKSAR. Updated Aug 12, 2021. Accessed Sep 21, 2021. https://www.censtatd.gov.hk/en/web_table.html?id=1B
14. Kwan MYW, Chua GT, Chow CB, et al. mRNA COVID vaccine and myocarditis in adolescents. *Hong Kong Med J*. Oct 2021;27(5):326-327. doi:10.12809/hkmj215120
15. Shay DK, Shimabukuro TT, DeStefano F. Myocarditis Occurring After Immunization With mRNA-Based COVID-19 Vaccines. *JAMA Cardiology*. 2021;doi:10.1001/jamacardio.2021.2821

16. Abu Mouch S, Roguin A, Hellou E, et al. Myocarditis following COVID-19 mRNA vaccination. *Vaccine*. Jun 29 2021;39(29):3790-3793. doi:10.1016/j.vaccine.2021.05.087
17. Simone A, Herald J, Chen A, et al. Acute Myocarditis Following COVID-19 mRNA Vaccination in Adults Aged 18 Years or Older. *JAMA Internal Medicine*. 2021;doi:10.1001/jamainternmed.2021.5511
18. Li C, Chen Y, Zhao Y, et al. Intravenous injection of COVID-19 mRNA vaccine can induce acute myopericarditis in mouse model. *Clin Infect Dis*. Aug 18 2021;doi:10.1093/cid/ciab707
19. Diaz GA, Parsons GT, Gering SK, Meier AR, Hutchinson IV, Robicsek A. Myocarditis and Pericarditis After Vaccination for COVID-19. *JAMA*. 2021;doi:10.1001/jama.2021.13443
20. Starekova J, Bluemke DA, Bradham WS, et al. Evaluation for Myocarditis in Competitive Student Athletes Recovering From Coronavirus Disease 2019 With Cardiac Magnetic Resonance Imaging. *JAMA Cardiol*. Jan 14 2021;doi:10.1001/jamacardio.2020.7444
21. Ghafoor K, Ahmed A, Abbas M. Fulminant Myocarditis With ST Elevation and Cardiogenic Shock in a SARS-CoV-2 Patient. *Cureus*. Jul 2021;13(7):e16149. doi:10.7759/cureus.16149
22. Małek Ł A, Marczak M, Miłosz-Wieczorek B, et al. Cardiac involvement in consecutive elite athletes recovered from Covid-19: A magnetic resonance study. *J Magn Reson Imaging*. Jun 2021;53(6):1723-1729. doi:10.1002/jmri.27513
23. Chodick G, Tene L, Patalon T, et al. The effectiveness of the first dose of BNT162b2 vaccine in reducing SARS-CoV-2 infection 13-24 days after immunization: real-world evidence. *medRxiv*. 2021:2021.01.27.21250612. doi:10.1101/2021.01.27.21250612
24. Latest situation of cases of COVID-19 (as of 18 November 2021) Centre for Health Protection, HKSAR. Accessed Nov 19, 2021. https://www.chp.gov.hk/files/pdf/local_situation_covid19_en.pdf
25. Scientific Committees under CHP issue consensus interim recommendations on COVID-19 vaccination. The Government of the Hong Kong Special Administrative Region. Accessed Sep 19, 2021. <https://www.info.gov.hk/gia/general/202109/15/P2021091501003.htm>

Table 1. Clinical characteristics of adolescents with myocarditis/pericarditis following Comirnaty vaccination in Hong Kong

N o	Sex/ Age at prese ntati on (year s)	Pre se nt aft er firs t or sec on d do se	No. of day s aft er rec eivi ng the last dos e	Symptoms	Peak Troponin Levels (hsTnT/h sTnI /TnT) (ng/L)#	Most Significant ECG Changes	ECHO	MRI findings	Final Diagnosis (Level of Certainty*)
1	M/15 .66	Sec on d	2	Chest pain headache	TnT 793 hsTnI 2506 (elevated)	STE in II,III, aVF, V3-V5	Normal	Patchy edema; diffuse EGE; patchy pericardial and subepicardial LGE; normal ECV	Perimyoca rditis (Definite)
2	M/14 .52	Sec on d	1	Chest pain fever	TnT 646 hsTnI 6342 (elevated)	TWI and biphasic T waves in III, aVF, V4-V6	Normal	Borderline LV function; Elevated T1 and T2 mapping values and ECV; presence of LGE	Myocarditi s (Definite)
3	M/13 .53	Sec on d	2	Dizziness, SOB, chest pain	TnT 1749 (elevated)	STE in V2-V6 TWI in aVL; biphasic Ts in V3-V4	Normal	Elevated T1 and T2 mapping values and ECV; pericardium and subepicardial muscles LGE and T2 hyperintensity	Perimyoca rditis (Definite)

4	M/13 .05	First	2	Chest pain	TnT 302 (elevated)	STE in V3-V6; TWI in III; STD in aVR	Normal	Elevated T1 and T2 mapping values and ECV; pericardial LGE extending to subepicardial region	Perimyocarditis (Definite)
5	M/14 .34	Second	1	Chest pain	TnT 993 (elevated)	STE in V3-V5; TWI in I, aVL; biphasic T waves in V3- V6	Normal	Borderline LV function; subepicardial LGE; elevated T1 and T2 mapping values and ECV; hyperintense pericardium	Perimyocarditis (Definite)
6	M/16 .99	Second	3	Chest pain	TnT 948 (elevated)	STE in V2-V6	Mildly impaired LV global longitudinal strain	Borderline LV function, small pericardial effusion; elevated ECV, T1 and T2 mapping values; patchy LGE	Perimyocarditis (Definite)
7	M/15 .22	Second	2	Chest pain	hsTnI 11415 (elevated)	Normal	Normal	Elevated T1 and T2 mapping values; presence of patchy EGE; normal ECV; subepicardial LGE	Myocarditis (Definite)
8	M/15 .32	Second	2	Chest pain fever	hsTnI 16806 (elevated)	STD and TWI in V1-2; STE in lead II, III, aVF; ST/T wave abnormality in II, III, aVF, V4-V6	Borderline LV function (LVFS 28%), minimal pericardial effusion	Mild increase STIR signal; faint patchy LGE; trace pericardial effusion	Perimyocarditis (Definite)
9	M/17 .14	First	1	Chest pain	hsTnI 19110 (elevated)	STE in II, III, aVF, V4-6	Tiny rim of pericardial effusion	Elevated T1 and T2 mapping values and ECV; no definite EGE; LGE present; patchy pericardial enhancement	Perimyocarditis (Definite)

10	F/14.07	Second	3	Chest discomfort, transient SOB	hsTnI 54.9 (elevated)	STE in V4-V5	Normal	Elevated T1 and T2 mapping values and ECV; LGE and pericardial enhancement	Perimyocarditis (Definite)
11	M/13.75	Second	2	Chest pain palpitation, fever	hsTnI 6254 (elevated)	Sinus tach; STE in II, III, aVF, V3-5	Normal	Elevated T1 and T2 mapping values and ECV; presence of LGE	Myocarditis (Definite)
12	M/12.74	Second	1	Chest pain, palpitations dizziness	hsTnI 14766 (elevated)	STD in aVR and V1; STE I-III, aVF, V4-6	Thin rim of pericardial effusion, hyperechoic pericardium	Elevated T1 mapping value; presence of myocardial edema with increased T2W signal	Perimyocarditis (Definite)
13	F/12.97	Second	1	Chest pain, fever, headache, palpitations, subjective SOB	hsTnI 2309 (elevated)	Normal	Normal	Elevated T1 and T2 mapping values and ECV; pericardial and subepicardial LGE; small pericardial effusion	Perimyocarditis (Definite)
14	M/17.85	Second	3	Chest pain	hsTnI 30267 (elevated)	STE in I, II, aVF, V4-6, STD in aVR, V1-V2; TWI in III; biphasic Ts in V3-V5	Borderline contractility	Elevated T1 and T2 mapping values and ECV; subepicardial and mid-wall LGE; small pericardial effusion	Perimyocarditis (Definite)
15	M/14.99	Second	1	Fever, chest pain, palpitation, SOB, dizziness	TnT 323 (elevated)	STE in V3-5 TWI in aVR and V1	Normal	T2W hyperintensity within myocardium; regional LGE; 5mm pericardial effusion	Perimyocarditis (Definite)
16	M/16.88	Second	4	Chest pain, SOB	hsTnT <14 (normal)	STE in V2 and V4	Increased echogenicity over LV free wall	Not done	Pericarditis (Definite)

17	M/17	Second	2	Chest pain, fever, palpitation	hsTnI 767 (elevated)	STE in II, III, aVF, V3-6	Normal	T2W hyperintense myocardial edema at mid and apex of LV	Myocarditis (Definite)
18	M/14	First	3	Chest pain, fever	hsTnI 184 (elevated)	STD in II, III, aVF	Normal	T2W hyperintense myocardial edema at basal lateral and basal septal segments of LV	Myocarditis (Definite)
19	M/15	Second	2	Chest discomfort, palpitation	hsTnI 3561 (elevated)	STE in II, aVF, V4-V6	Normal	T2W hyperintense myocardial edema with LGE at apical lateral segment and subepicardial region	Myocarditis (Definite)
20	M/14	Second	2	Chest pain	TnT 1058 (elevated)	STE in V4-V6; TWI in III, aVF	Normal	Mild T2W hyperintense signals and increased T2 mapping value at inferolateral LV wall	Myocarditis (Definite)
21	M/15	Second	2	Chest pain	hsTnI 263 (elevated)	STE II,III,aVF, V4-V6	Normal	Mild subepicardial basal to mid-ventricular lateral wall LGE and elevated T1 mapping value	Myocarditis (Definite)
22	M/15	Second	1	Chest pain, palpitations	hsTnI 2210 TnT 283 (elevated)	STE V2-6	Normal	Generalized myocardial hyperintensity in TIRM sequence; presence of hyperemia; subepicardial LGE; small pericardial effusion	Perimyocarditis (Definite)
23	F/16.89	First	2	palpitation, near syncope, nausea, vomiting	hsTnT 30 (elevated)	Normal	Normal	LV myocardium diffuse increased T2 signal; patchy early Gd enhancement	Myocarditis (Definite)
24	M/16	Second	2	chest pain, headache, dizziness	TnT 669 (elevated)	Normal	Borderline LV function (LVFS 29.1%)	Normal	Myocarditis (Definite)
25	M/14	Second	2	Chest pain, palpitation	hsTnT <14 (normal)	STE in I,II, V2-6, and STD in aVR	Prominent pericardial echogenicity	Normal	Pericarditis (Definite)

26	M/14 .18	First	2	Chest pain	hsTnI 513 (elevated)	Normal	Normal	Equivocal myocardial edema due to motion artefacts	Myocarditis (Probable)
27	F/15. 25	Second	6	Chest pain	hsTnI 77 (elevated)	STE V2-3; biphasic Ts in V3	Normal	Normal	Myocarditis (Probable)
28	M/14 .31	Second	14	Chest discomfort, transient SOB, headache, dizziness	hsTnI 201 (elevated)	TWI and ST depression in II, III, aVF; biphasic Ts in V3-V5	Normal	Normal	Myocarditis (Probable)
29	M/17 .87	First	2	Chest pain	hsTnI 29.2 (elevated)	STE in II, V3- 6	Normal	Normal	Myocarditis (Probable)
30	M/17 .64	Second	2	Chest pain, fever headache	hsTnI 4874 (elevated)	STE in V2-6, TWI in aVF/III; biphasic Ts in II, aVF, V4- V6	Normal	Normal	Myocarditis (Probable)
31	M/12 .85	Second	2	Chest pain, vomiting, SOB	hsTnT 39	Sinus tachycardia; STE in II, III, aVF; V2-V6	Normal	Global hyperintensity in myocardium in T2W images with hyperintensity in early post-Gd images but no LGE. Suspicious of myocarditis	Myocarditis (Probable)
32	M/15 .79	Second	10	chest pain, dizziness, near syncope	hsTnT 25 (elevated)	Normal	Normal	Normal	Myocarditis (Probable)

3 3	M/16 .76	Sec on d	2	Fever chest discomfort, palpitation, transient SOB	hsTnT <14 (normal)	STE in V2-V6	Normal	Normal	Myocarditi s (Possible – elevated CRP)
3 4 ^	M/15 .07	Sec on d	25	Chest Pain	TnT 269 hsTnI 3850	STE in V2-V6	Mild pericardial and LV free wall echogenicity	Not done	Perimyoca rditis (Definite)
3 5 ^	F/12. 78	Sec on d	26	Vomiting, palpitation, reduced exercise tolerance	hsTnI 566	STE in II, V2- V5; STD in aVR; TWI in aVL; Q waves in I and aVL	Hyperechoic pericardium	Elevated T1 mapping values; subepicardial LGE	Perimyoca rditis (Definite)

*Brighton Collaboration Myocarditis Case Definition Level of Certainty (LOC) Classification

Elevated troponin level based on reference values provided by each laboratory. Subjects with two different troponin measures were because of transferral to another hospital.

^ Cases 34 and 35 presented >14 days after receiving the second doses, therefore they were only included in the sensitivity analyses (supplementary table 4 – 8)

CRP – C reactive protein; ECG – electrocardiogram; ECHO – echocardiogram; ECV – extracellular volume; EGE – early gadolinium enhancement; Gd – Gadolinium; hsTnI – high-sensitivity troponin I; hsTnT – high-sensitivity troponin T; LGE – late gadolinium enhancement; LV – left ventricle; LVFS – left ventricle fractional shortening; SOB – shortness of breath; STD – ST depression; STE – ST elevation; STIR – short tau inversion recovery; T2W – T2-weighted; TWI – T wave inversion; TnT – troponin T

Table 2. Incidence rate differences of myocarditis/pericarditis cases following Comirnaty vaccination stratified by sex and compared to background rate in 2020

	Incidence rate (per 100,000 person-14 days, 95% CI)	Background incidence rate in 2020 [†] (per 100,000 person-14 days, 95% CI)	Incidence rate difference (per 100,000 person-14 days, 95% CI)
Comirnaty			
Total	18.52 (11.67-29.01)	0.11 (0.01-20.36)	18.41 (9.95-26.87)
Male	32.29 (22.78-45.4)	0.21 (0.01-10.34)	32.08 (20.91-43.25)
Female	4.53 (1.76-11.11)	0	-

[†]The background incidence rates were calculated using the reporting period (14 June – 4 September) in 2020 and truncated to incidence rate per 14 days. Values in **bold** represent a statistically significant difference (P<0.05).

Table 3. Incidence rate differences of myocarditis/pericarditis cases following the first and second doses of Comirnaty vaccination stratified by sex and compared to background rate in 2020

	Incidence rate (per 100,000 person-14 days, 95% CI)	Background incidence rate in 2020† (per 100,000 person-14 days, 95% CI)	Incidence rate difference (per 100,000 person-14 days, 95% CI)
First dose of Comirnaty			
Total	3.37 (1.12-9.51)	0.11 (0.01-20.36)	3.26 (-0.40-6.92)
Male	5.57 (2.38-12.53)	0.21 (0.01-10.34)	5.36 (0.65-10.07)
Females	1.13(0.16-6.58)	0	-
Second dose of Comirnaty			
Total	21.22 (13.78-32.28)	0.11 (0.01-20.36)	21.11 (12.06-30.16)
Male	37.32 (26.98-51.25)	0.21 (0.01-10.34)	37.11 (25.10-49.12)
Female	4.77 (1.90-11.44)	0	-

†The background incidence rates were calculated using the reporting period (14 June – 4 September) in 2020 and truncated to incidence rate per 14 days. Values in **bold** represent a statistically significant difference (P<0.05).